

5. PROPAGATION OF VARIANCES APPLIED TO LEAST SQUARES

ADJUSTMENT OF INDIRECT OBSERVATIONS

A most important outcome of a least squares adjustment is that estimates of the precisions of the quantities sought, the elements of \mathbf{x} , the unknowns or the parameters, are easily obtained from the matrix equations of the solution. Application of the Law of Propagation of Variances demonstrates that \mathbf{N}^{-1} , the inverse of the normal equation coefficient matrix is equal to the cofactor matrix \mathbf{Q}_{xx} that contains estimates of the variances and covariances of the elements of \mathbf{x} . In addition, estimates of the precisions of the residuals and adjusted observations may be obtained. This most useful outcome enables a statistical analysis of the results of a least squares adjustment and provides the practitioner with a degree of confidence in the results.

5.1. Cofactor matrices for adjustment of indirect observations

The observation equations for adjustment of indirect observations is given by

$$\mathbf{v} + \mathbf{B}\mathbf{x} = \mathbf{f} \quad (5.1)$$

\mathbf{f} is an $(n, 1)$ vector of numeric terms derived from the $(n, 1)$ vector of observations \mathbf{l} and the $(n, 1)$ vector of constants \mathbf{d} as

$$\mathbf{f} = \mathbf{d} - \mathbf{l} \quad (5.2)$$

Associated with the vector of observations \mathbf{l} is a variance-covariance matrix Σ_{ll} as well as a cofactor matrix \mathbf{Q}_{ll} and a weight matrix $\mathbf{W}_{ll} = \mathbf{Q}_{ll}^{-1}$. Remember that in most practical applications of least squares, the matrix Σ_{ll} is unknown, but estimated *a priori* by \mathbf{Q}_{ll} that contains estimates of the variances and covariances and $\Sigma_{ll} = \sigma_0^2 \mathbf{Q}_{ll}$ where σ_0^2 is the reference variance or variance factor.

Note: In the derivations that follow, the subscript "ll" is dropped from \mathbf{Q}_{ll} and \mathbf{W}_{ll}

If equation (5.2) is written as

$$\mathbf{f} = (-\mathbf{I})\mathbf{l} + \mathbf{d} \quad (5.3)$$

then (5.3) is in a form suitable for employing the Law of Propagation of Variances developed in Chapter 3; i.e., if $\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$ and \mathbf{y} and \mathbf{x} are random variables linearly related and \mathbf{b} is a vector of constants then $\mathbf{Q}_{yy} = \mathbf{A}\mathbf{Q}_{xx}\mathbf{A}^T$. Hence, the cofactor matrix of the numeric terms \mathbf{f} is

$$\mathbf{Q}_{ff} = (-\mathbf{I})\mathbf{Q}(-\mathbf{I})^T = \mathbf{Q}$$

Thus the cofactor matrix of \mathbf{f} is also the *a priori* cofactor matrix of the observations \mathbf{l} .

The solution "steps" in the least squares adjustment of indirect observations are set out Chapter 2 and restated as

$$\begin{aligned}\mathbf{N} &= \mathbf{B}^T \mathbf{W} \mathbf{B} \\ \mathbf{t} &= \mathbf{B}^T \mathbf{W} \mathbf{f} \\ \mathbf{x} &= \mathbf{N}^{-1} \mathbf{t} \\ \mathbf{v} &= \mathbf{f} - \mathbf{B}\mathbf{x} \\ \hat{\mathbf{l}} &= \mathbf{l} + \mathbf{v}\end{aligned}$$

To apply the Law of Propagation of Variances, these equations may be re-arranged in the form $\mathbf{y} = \mathbf{A}\mathbf{x} + \mathbf{b}$ where the terms in parenthesis () constitute the \mathbf{A} matrix.

$$\mathbf{t} = (\mathbf{B}^T \mathbf{W}) \mathbf{f} \quad (5.4)$$

$$\mathbf{x} = (\mathbf{N}^{-1}) \mathbf{t} \quad (5.5)$$

$$\begin{aligned}\mathbf{v} &= \mathbf{f} - \mathbf{B}\mathbf{x} \\ &= \mathbf{f} - \mathbf{B}\mathbf{N}^{-1}\mathbf{t} \\ &= \mathbf{f} - \mathbf{B}\mathbf{N}^{-1}\mathbf{B}^T \mathbf{W} \mathbf{f} \\ &= (\mathbf{I} - \mathbf{B}\mathbf{N}^{-1}\mathbf{B}^T \mathbf{W}) \mathbf{f}\end{aligned} \quad (5.6)$$

$$\begin{aligned}\hat{\mathbf{l}} &= \mathbf{l} + \mathbf{v} \\ &= \mathbf{l} + \mathbf{f} - \mathbf{B}\mathbf{x} \\ &= \mathbf{d} - \mathbf{B}\mathbf{x} \\ &= (-\mathbf{B})\mathbf{x} + \mathbf{d}\end{aligned} \quad (5.7)$$

Applying the Law of Propagation of Variances to equations (5.4) to (5.7) gives the following cofactor matrices

$$\mathbf{Q}_{tt} = (\mathbf{B}^T \mathbf{W}) \mathbf{Q}_{ff} (\mathbf{B}^T \mathbf{W})^T = \mathbf{N} \quad (5.8)$$

$$\mathbf{Q}_{xx} = (\mathbf{N}^{-1}) \mathbf{Q}_{tt} (\mathbf{N}^{-1})^T = \mathbf{N}^{-1} \quad (5.9)$$

$$\begin{aligned} \mathbf{Q}_{vv} &= (\mathbf{I} - \mathbf{B} \mathbf{N}^{-1} \mathbf{B}^T \mathbf{W}) \mathbf{Q}_{ff} (\mathbf{I} - \mathbf{B} \mathbf{N}^{-1} \mathbf{B}^T \mathbf{W})^T \\ &= \mathbf{Q} - \mathbf{B} \mathbf{N}^{-1} \mathbf{B}^T \end{aligned} \quad (5.10)$$

$$\begin{aligned} \mathbf{Q}_{\hat{\hat{t}}} &= (-\mathbf{B}) \mathbf{Q} (-\mathbf{B})^T \\ &= \mathbf{B} \mathbf{N}^{-1} \mathbf{B}^T \\ &= \mathbf{Q} - \mathbf{Q}_{vv} \end{aligned} \quad (5.11)$$

Variance-covariance matrices for \mathbf{t} , \mathbf{x} , \mathbf{v} and $\hat{\hat{\mathbf{t}}}$ are obtained by multiplying the cofactor matrix by the variance factor σ_0^2 .

5.2. Calculation of the quadratic form $\mathbf{v}^T \mathbf{W} \mathbf{v}$

The *a priori* estimate of the variance factor may be computed from

$$\hat{\sigma}_0^2 = \frac{\mathbf{v}^T \mathbf{W} \mathbf{v}}{r} \quad (5.12)$$

where $\mathbf{v}^T \mathbf{W} \mathbf{v}$ is the quadratic form, and

$r = n - u$ is the degrees of freedom where n is the number of observations and u is the number of unknown parameters. r is also known as the number of redundancies.

A derivation of equation (5.12) is given below. The quadratic form $\mathbf{v}^T \mathbf{W} \mathbf{v}$ may be computed in the following manner.

Remembering, for the method of indirect observations, the following matrix equations

$$\begin{aligned} \mathbf{N} &= \mathbf{B}^T \mathbf{W} \mathbf{B} \\ \mathbf{t} &= \mathbf{B}^T \mathbf{W} \mathbf{f} \\ \mathbf{x} &= \mathbf{N}^{-1} \mathbf{t} \\ \mathbf{v} &= \mathbf{f} - \mathbf{B} \mathbf{x} \end{aligned}$$

then

$$\begin{aligned}
\mathbf{v}^T \mathbf{W} \mathbf{v} &= (\mathbf{f} - \mathbf{B} \mathbf{x})^T \mathbf{W} (\mathbf{f} - \mathbf{B} \mathbf{x}) \\
&= (\mathbf{f}^T - \mathbf{x}^T \mathbf{B}^T) \mathbf{W} (\mathbf{f} - \mathbf{B} \mathbf{x}) \\
&= (\mathbf{f}^T \mathbf{W} - \mathbf{x}^T \mathbf{B}^T \mathbf{W}) (\mathbf{f} - \mathbf{B} \mathbf{x}) \\
&= \mathbf{f}^T \mathbf{W} \mathbf{f} - \mathbf{f}^T \mathbf{W} \mathbf{B} \mathbf{x} - \mathbf{x}^T \mathbf{B}^T \mathbf{W} \mathbf{f} + \mathbf{x}^T \mathbf{B}^T \mathbf{W} \mathbf{B} \mathbf{x} \\
&= \mathbf{f}^T \mathbf{W} \mathbf{f} - 2\mathbf{f}^T \mathbf{W} \mathbf{B} \mathbf{x} + \mathbf{x}^T \mathbf{B}^T \mathbf{W} \mathbf{B} \mathbf{x} \\
&= \mathbf{f}^T \mathbf{W} \mathbf{f} - 2\mathbf{t}^T \mathbf{x} + \mathbf{x}^T \mathbf{N} \mathbf{x} \\
&= \mathbf{f}^T \mathbf{W} \mathbf{f} - 2\mathbf{x}^T \mathbf{t} + \mathbf{x}^T \mathbf{t}
\end{aligned}$$

and

$$\mathbf{v}^T \mathbf{W} \mathbf{v} = \mathbf{f}^T \mathbf{W} \mathbf{f} - \mathbf{x}^T \mathbf{t} \quad (5.13)$$

5.3. Calculation of the Estimate of the Variance Factor $\hat{\sigma}_0^2$

The variance-covariance matrices of residuals Σ_{vv} , adjusted observations $\Sigma_{\hat{y}}$ and computed parameters Σ_{xx} are calculated from the general relationship

$$\Sigma = \sigma_0^2 \mathbf{Q} \quad (5.14)$$

Cofactor matrices \mathbf{Q}_{vv} , \mathbf{Q}_{xx} and $\mathbf{Q}_{\hat{y}}$ are computed from equations (5.9) to (5.11) and so it remains to determine an estimate of the variance factor $\hat{\sigma}_0^2$.

The development of a matrix expression for computing $\hat{\sigma}_0^2$ is set out below and follows Mikhail (1976, pp.285-288). Some preliminary relationships will be useful.

1. If \mathbf{A} is an (n, n) square matrix, the sum of its diagonal elements is a scalar quantity called the trace of \mathbf{A} and denoted by $tr(\mathbf{A})$. The following relationships are useful

$$tr(\mathbf{A} + \mathbf{B}) = tr(\mathbf{A}) + tr(\mathbf{B}) \quad \text{for } \mathbf{A} \text{ and } \mathbf{B} \text{ of same order} \quad (5.15)$$

$$tr(\mathbf{A}^T) = tr(\mathbf{A}) \quad (5.16)$$

and for the quadratic form $\mathbf{x}^T \mathbf{A} \mathbf{x}$ where \mathbf{A} is symmetric

$$\mathbf{x}^T \mathbf{A} \mathbf{x} = tr(\mathbf{x} \mathbf{x}^T \mathbf{A}) \quad (5.17)$$

2. The variance-covariance matrix Σ_{xx} given by equation (3.21) can be expressed in the following manner, remembering that \mathbf{x} is a vector of random variables and \mathbf{m}_x is a vector of means.

$$\begin{aligned}\Sigma_{xx} &= E\{(\mathbf{x} - \mathbf{m}_x)(\mathbf{x} - \mathbf{m}_x)^T\} \\ &= E\{(\mathbf{x} - \mathbf{m}_x)(\mathbf{x}^T - \mathbf{m}_x^T)\} \\ &= E\{\mathbf{xx}^T - \mathbf{xm}_x^T - \mathbf{m}_x\mathbf{x}^T + \mathbf{m}_x\mathbf{m}_x^T\} \\ &= E\{\mathbf{xx}^T\} - E\{\mathbf{xm}_x^T\} - E\{\mathbf{m}_x\mathbf{x}^T\} + E\{\mathbf{m}_x\mathbf{m}_x^T\} \\ &= E\{\mathbf{xx}^T\} - E\{\mathbf{x}\}\mathbf{m}_x^T - \mathbf{m}_xE\{\mathbf{x}^T\} + \mathbf{m}_x\mathbf{m}_x^T\end{aligned}$$

Now from equation (3.18) $\mathbf{m}_x = E\{\mathbf{x}\}$ hence

$$\begin{aligned}\Sigma_{xx} &= E\{\mathbf{xx}^T\} - \mathbf{m}_x\mathbf{m}_x^T - \mathbf{m}_x\mathbf{m}_x^T + \mathbf{m}_x\mathbf{m}_x^T \\ &= E\{\mathbf{xx}^T\} - \mathbf{m}_x\mathbf{m}_x^T\end{aligned}\quad (5.18)$$

or
$$E\{\mathbf{xx}^T\} = \Sigma_{xx} + \mathbf{m}_x\mathbf{m}_x^T \quad (5.19)$$

3. The expected value of the residuals is zero, i.e.,

$$E\{\mathbf{v}\} = \mathbf{m}_v = \mathbf{0} \quad (5.20)$$

4. By definition (see Chapter 2) the weight matrix \mathbf{W} , the cofactor matrix \mathbf{Q} and the variance-covariance matrix Σ are related by

$$\mathbf{W} = \mathbf{Q}^{-1} = \sigma_0^2 \Sigma^{-1} \quad (5.21)$$

Now, for the least squares adjustment of indirect observations the following relationships are recalled

$$\begin{aligned}\mathbf{v} + \mathbf{Bx} &= \mathbf{f}, \quad \mathbf{N} = \mathbf{B}^T \mathbf{W} \mathbf{B}, \quad \mathbf{t} = \mathbf{B}^T \mathbf{W} \mathbf{f} \\ \mathbf{Q}_{ff} &= \mathbf{Q}, \quad \mathbf{Q}_{tt} = \mathbf{N}, \quad \mathbf{Q}_{xx} = \mathbf{N}^{-1}\end{aligned}$$

Bearing in mind equation (5.21), the following relationships may be introduced

$$\Sigma^{-1} = \frac{1}{\sigma_0^2} \mathbf{W}, \quad \mathbf{M} = \mathbf{B}^T \Sigma^{-1} \mathbf{B}$$

and from these follow

$$\Sigma_{ff} = \Sigma, \quad \Sigma_{tt} = \sigma_0^4 \mathbf{M}, \quad \Sigma_{xx} = \mathbf{M}^{-1}$$

In addition, the expectation of the vector \mathbf{f} is the mean \mathbf{m}_f and so we may write

$$\mathbf{m}_f = E\{\mathbf{f}\} = E\{\mathbf{v} + \mathbf{B}\mathbf{x}\} = E\{\mathbf{v}\} + \mathbf{B}E\{\mathbf{x}\}$$

Now since $E\{\mathbf{x}\} = \mathbf{m}_x$ and $E\{\mathbf{v}\} = \mathbf{0}$

$$\mathbf{m}_f = \mathbf{B}\mathbf{m}_x \quad (5.22)$$

Now the quadratic form

$$\mathbf{v}^T \mathbf{W}\mathbf{v} = \sigma_0^2 (\mathbf{v}^T \Sigma^{-1} \mathbf{v}) \quad (5.23)$$

and from equation (5.13)

$$\begin{aligned} \mathbf{v}^T \mathbf{W}\mathbf{v} &= \mathbf{f}^T \mathbf{W}\mathbf{f} - \mathbf{x}^T \mathbf{t} \\ &= \mathbf{f}^T \mathbf{W}\mathbf{f} - \mathbf{x}^T \mathbf{N}\mathbf{x} \end{aligned}$$

Using the relationships above

$$\mathbf{v}^T \Sigma^{-1} \mathbf{v} = \mathbf{f}^T \Sigma^{-1} \mathbf{f} - \mathbf{x}^T \mathbf{M}\mathbf{x}$$

Now the expected value of this quadratic form is

$$\begin{aligned} E\{\mathbf{v}^T \Sigma^{-1} \mathbf{v}\} &= E\{\mathbf{f}^T \Sigma^{-1} \mathbf{f} - \mathbf{x}^T \mathbf{M}\mathbf{x}\} \\ &= E\{\mathbf{f}^T \Sigma^{-1} \mathbf{f}\} - E\{\mathbf{x}^T \mathbf{M}\mathbf{x}\} \end{aligned}$$

Recognising that the terms on the right-hand-side are both quadratic forms, equation (5.17) can be used to give

$$\begin{aligned} E\{\mathbf{v}^T \Sigma^{-1} \mathbf{v}\} &= E\{tr(\mathbf{f}\mathbf{f}^T \Sigma^{-1})\} - E\{tr(\mathbf{x}\mathbf{x}^T \mathbf{M})\} \\ &= tr(E\{\mathbf{f}\mathbf{f}^T \Sigma^{-1}\}) - tr(E\{\mathbf{x}\mathbf{x}^T \mathbf{M}\}) \\ &= tr(E\{\mathbf{f}\mathbf{f}^T\} \Sigma^{-1}) - tr(E\{\mathbf{x}\mathbf{x}^T\} \mathbf{M}) \end{aligned}$$

Now using equation (5.19)

$$\begin{aligned} E\{\mathbf{v}^T \Sigma^{-1} \mathbf{v}\} &= tr\left(\left[\Sigma_{ff} + \mathbf{m}_f \mathbf{m}_f^T\right] \Sigma^{-1}\right) - tr\left(\left[\Sigma_{xx} + \mathbf{m}_x \mathbf{m}_x^T\right] \mathbf{M}\right) \\ &= tr\left(\mathbf{I}_{nn} + \mathbf{m}_f \mathbf{m}_f^T \Sigma^{-1}\right) - tr\left(\mathbf{I}_{uu} + \mathbf{m}_x \mathbf{m}_x^T \mathbf{M}\right) \\ &= tr\left(\mathbf{I}_{nn} - \mathbf{I}_{uu}\right) - tr\left(\mathbf{m}_f \mathbf{m}_f^T \Sigma^{-1} + \mathbf{m}_x \mathbf{m}_x^T \mathbf{M}\right) \\ &= (n - u) - \mathbf{m}_f^T \Sigma^{-1} \mathbf{m}_f + \mathbf{m}_x^T \mathbf{M} \mathbf{m}_x \end{aligned}$$

From equation (5.22) $\mathbf{m}_f = \mathbf{Bm}_x$ hence using the rule for matrix transpose

$\mathbf{m}_f^T = (\mathbf{Bm}_x)^T = \mathbf{m}_x^T \mathbf{B}^T$, then

$$\begin{aligned} E\{\mathbf{v}^T \boldsymbol{\Sigma}^{-1} \mathbf{v}\} &= (n - u) - \mathbf{m}_x^T \mathbf{B}^T \boldsymbol{\Sigma}^{-1} \mathbf{Bm}_x + \mathbf{m}_x^T \mathbf{Mm}_x \\ &= (n - u) - \mathbf{m}_x^T \mathbf{Mm}_x + \mathbf{m}_x^T \mathbf{Mm}_x \\ &= (n - u) \end{aligned}$$

Thus according to equation (5.23) and the expression above

$$\begin{aligned} E\{\mathbf{v}^T \mathbf{Wv}\} &= \sigma_0^2 E\{\mathbf{v}^T \boldsymbol{\Sigma}^{-1} \mathbf{v}\} \\ &= \sigma_0^2 (n - u) \end{aligned}$$

from which follows

$$\sigma_0^2 = \frac{E\{\mathbf{v}^T \mathbf{Wv}\}}{n - u}$$

Consequently, an unbiased estimate of the variance factor $\hat{\sigma}_0^2$ can be computed from

$$\hat{\sigma}_0^2 = \frac{\mathbf{v}^T \mathbf{Wv}}{n - u} = \frac{\mathbf{v}^T \mathbf{Wv}}{r} \quad (5.24)$$

$r = n - u$ is the number of redundancies in the adjustment and is known as the degrees of freedom

Using equation (5.13) an unbiased estimate of the variance factor $\hat{\sigma}_0^2$ can be computed from

$$\hat{\sigma}_0^2 = \frac{\mathbf{f}^T \mathbf{Wf} - \mathbf{x}^T \mathbf{t}}{r} \quad (5.25)$$